



TITLE:

Light-dependent destabilization of PHL in Arabidopsis

AUTHOR(S):

Endo, Motomu; Kudo, Daiki; Koto, Tomoko;
Shimizu, Hanako; Araki, Takashi

CITATION:

Endo, Motomu ...[et al]. Light-dependent destabilization of PHL in Arabidopsis. Plant Signaling & Behavior 2014, 9(2): e28118.

ISSUE DATE:

2014-03-10

URL:

<http://hdl.handle.net/2433/187356>

RIGHT:

© 2014 Landes Bioscience; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。; This is not the published version. Please cite only the published version.

1 ***Short Communication***

2

3 **Title**

4 Light-dependent destabilization of PHL in Arabidopsis

5

6 **Authors**

7 Motomu Endo*, Daiki Kudo, Tomoko Koto, Hanako Shimizu, and Takashi Araki*

8

9 **Affiliation**

10 Division of Integrated Life Science, Graduate School of Biostudies, Kyoto University,

11 Kyoto 606-8501, Japan

12

13 **Key words**

14 Arabidopsis, flowering, florigen, phytochrome, photoperiod, FT, CO, phyB, PHL

15

16 ***Co-correspondence to:**

17 Motomu Endo or Takashi Araki; Division of Integrated Life Science, Graduate School

18 of Biostudies, Kyoto University, Kyoto 606-8501, Japan

19

20 Tel: +81-75-753-6140; Fax: +81-75-753-6470; Email: moendo@lif.kyoto-u.ac.jp (ME)

21 or taraqui@lif.kyoto-u.ac.jp (TA)

22

23 **Abbreviations**

24 Cauliflower Mosaic Virus, CaMV; CONSTANS, CO; CONSTITUTIVE

25 PHOTOMORPHOGENIC 1, COP1; FLOWERING LOCUS T, FT; long day, LD;

26 Phytochrome B, phyB; PHYTOCHROME-DEPENDENT LATE-FLOWERING, PHL;

27 short day, SD.

28

29 **Abstract**

30 Plants sense environmental stimuli such as light to regulate their flowering time. In

31 *Arabidopsis*, phytochrome B (phyB) is the major photoreceptor that perceives red and

32 far-red light, and destabilizes transcriptional regulator CONSTANS (CO) protein.

33 However the mechanism that links photoreceptor and CO protein degradation is largely

34 unknown. We recently showed that PHYTOCHROME-DEPENDENT

35 LATE-FLOWERING (PHL) protein inhibits phyB signaling through direct

36 protein-protein interaction. Here, we report that light exposure destabilizes PHL protein

as is the case with CO. Fluorescence from PHL-YFP fusion protein expressed under the control of Cauliflower Mosaic Virus (CaMV) 35S promoter (35S::PHL-YFP) almost disappeared after four-hour treatment of white light. Furthermore, the similar results were also obtained from the analysis of PHL-GUS fusion protein expressed by PHL promoter (PHL_{pro}::PHL-GUS *phl-1*). These results highlight the importance of post-transcriptional regulation in phyB-mediated flowering regulation and will give us hints how phyB regulates CO protein amount.

Text

Since light is one of the most important environmental signals in plants, various kinds of photoreceptors have evolved.¹ Among them, a red/far-red light photoreceptor phyB and several blue light photoreceptors regulate flowering time through modulating CO protein stability.²⁻⁴ However, how phyB regulates CO protein amount has remained unclear. Our recent work demonstrated that a novel protein, PHL, interacts with phyB *in vitro* and *in vivo*.⁵ Furthermore, two *phl* mutant alleles cause late-flowering phenotype under long day (LD) but not under short day (SD) conditions, suggesting that PHL regulates flowering in the photoperiod pathway. Consistent with the view, *FLOWERING LOCUS T* (*FT*) expression under LD condition was suppressed in the *phl*

mutant. These findings suggest that PHL have significant roles in flowering regulation by modulating phyB-signaling pathway. It was also demonstrated that the PHL could bridge interaction between phyB and CO protein in a red-light-dependent manner, implying that PHL protein may also undergo light-dependent destabilization, as does CO.²

Here, we report that PHL protein is destabilized in response to light exposure. We first established a transgenic line that expresses PHL fused to YFP, under the control of the CaMV 35S promoter in the wild-type background (35S::PHL-YFP). Since the 35S::PHL-YFP line produced a 2-fold elevated *PHL* mRNA level, the line was expected to have only slight side effects of exogenous PHL-YFP (data not shown). In consistent with the low expression level of PHL-YFP, the transgenic line showed no significant phenotype both under LD and SD conditions (Fig. 1A, B). To test the hypothesis that PHL is destabilized by light exposure, we observed fluorescence of PHL-YFP in dark- and light-grown seedlings. YFP fluorescence was observed in dark-grown seedlings, whereas significant fluorescence was not observed in light-grown seedlings (Fig. 1C). Since the CaMV 35S promoter are active both under light and dark conditions, posttranscriptional regulation of PHL by light is strongly suggested. We then performed time-course observation of the PHL-YFP fluorescence. Dark-grown plants were

73 transferred to continuous white light condition for 24 hours. The intensity of PHL-YFP
74 fluorescence was decreased in proportion to the time under continuous white light, and
75 no significant fluorescence was observed after four-hour exposure to light (Fig 1C).

76 To confirm these observations, we also employed the PHLpro:PHL-GUS *phl-1*,
77 which was used in our previous study.⁵ PHL protein amount in seedlings was examined
78 by staining for GUS (Fig. 2A). In consistent with the observation from PHL-YFP,
79 enough amount of PHL-GUS was detected in the dark grown seedlings, whereas
80 PHL-GUS accumulation was not detected in the light grown seedlings (Fig. 2A).
81 Kinetics of PHL-GUS protein degradation was also comparable to that of PHL-YFP
82 (Fig. 1C and Fig.2B). Furthermore, accumulation of PHL was observed only in
83 cotyledons even though the *PHL* mRNA expression has been detected in all organs
84 tested (Fig. 2A).⁵

85 Through the time-course observation of PHL protein expressed as fusion proteins,
86 we showed that PHL protein is destabilized by light exposure. Since phyB and PHL
87 interact directly in a red-light-dependent manner, it is likely that PHL is degraded in
88 response to red light. In support of this hypothesis, PHYTOCHROME INTERACTING
89 FACTORS (PIFs) and CO are also destabilized by red-light exposure, suggesting that
90 these proteins are destabilized in a similar mechanism.^{2,6-9} Previous studies

demonstrated that an E3 ubiquitin ligase, CONSTITUTIVE PHOTOMORPHOGENIC 1 (COP1), is involved in the destabilization process of CO and PIF1.^{7,10,11} Therefore, future study should involve the analysis of protein interaction between COP1 and PHL.

Interestingly, not only the PHLpro::PHL-GUS line but also the 35S::PHL-YFP line showed leaf-specific expression of PHL in the dark-grown seedlings, indicating the existence of an active destabilization mechanism of PHL presumably operating independently of phyB, although the biological meanings of the organ-specific degradation is unclear. Previous studies also demonstrated that phyB and CO regulate flowering by acting in leaves, supporting the existence of functional phyB-PHL-CO tripartite complex in leaves.^{5, 12,13}

In conclusion, our study provides a new insight into the phyB-mediated and phyB-independent protein degradation system(s). Together with our recent findings, it is suggested that destabilization of PHL is an important step to modulate phyB signaling in the photoperiod pathway. Therefore, elucidation of molecular mechanism of PHL protein destabilization will help to understand how phyB regulates flowering by modulating CO protein amount.

Acknowledgments

109 This work was in part supported by a JSPS KAKENHI grant 22770036 (to M.E.),
110 Grants-in-Aid for Scientific Research on Priority Areas 19060012 and 19060016 (to
111 T.A.), and a Grant-in-Aid for Scientific Research on Innovative Areas 25113005 (to
112 T.A.).

113

114 **References**

115 1. Kami C, Lorrain S, Hornitschek P, Fankhauser C. Light-regulated plant growth and
116 development. *Curr Top Dev Biol.* 2010;91:29-66.

117 2. Valverde F, Mouradov A, Soppe W, Ravenscroft D, Samach A, Coupland G.
118 Photoreceptor regulation of CONSTANS protein in photoperiodic flowering. *Science*
119 2004;303:1003-6.

120 3. Zuo Z, Liu H, Liu B, Liu X, Lin C. Blue light-dependent interaction of CRY2 with
121 SPA1 regulates COP1 activity and floral initiation in Arabidopsis. *Curr Biol.* 2011
122 24;21:841-7.

123 4. Song YH, Smith RW, To BJ, Millar AJ, Imaizumi T. FKF1 conveys timing
124 information for CONSTANS stabilization in photoperiodic flowering. *Science* 2012
125 25;336:1045-9.

126 5. Endo M, Tanigawa Y, Murakami T, Araki T, Nagatani A.

- 127 PHYTOCHROME-DEPENDENT LATE-FLOWERING accelerates flowering through
- 128 physical interactions with phytochrome B and CONSTANS. Proc Natl Acad Sci USA
- 129 2013;110:18017-22.
- 130 6. Park E, Kim J, Lee Y, Shin J, Oh E, Chung WI, Liu JR, Choi G. Degradation of
- 131 phytochrome interacting factor 3 in phytochrome-mediated light signaling. Plant Cell
- 132 Physiol. 2004;45:968-75.
- 133 7. Shen H, Moon J, Huq E. PIF1 is regulated by light-mediated degradation through the
- 134 ubiquitin-26S proteasome pathway to optimize photomorphogenesis of seedlings in
- 135 *Arabidopsis*. Plant J. 2005;44:1023-35.
- 136 8. Oh E, Yamaguchi S, Kamiya Y, Bae G, Chung WI, Choi G. Light activates the
- 137 degradation of PIL5 protein to promote seed germination through gibberellin in
- 138 *Arabidopsis*. Plant J. 2006;47:124-39.
- 139 9. Nozue K, Covington MF, Duek PD, Lorrain S, Fankhauser C, Harmer SL, Maloof JN.
- 140 Rhythmic growth explained by coincidence between internal and external cues. Nature
- 141 2007;448:358-61.
- 142 10. Jang S, Marchal V, Panigrahi KC, Wenkel S, Soppe W, Deng XW, Valverde F,
- 143 Coupland G. *Arabidopsis* COP1 shapes the temporal pattern of CO accumulation
- 144 conferring a photoperiodic flowering response. EMBO J. 2008;27:1277-88.

145 11. Laubinger S, Marchal V, Le Gourrierc J, Wenkel S, Adrian J, Jang S, Kulajta C,
146 Braun H, Coupland G, Hoecker U. *Arabidopsis* SPA proteins regulate photoperiodic
147 flowering and interact with the floral inducer CONSTANS to regulate its stability.
148 Development. 2006;133:3213-22.

149 12. Zeevaart, J.A.D. Physiology of flower formation. Annu. Rev. Plant Physiol.
150 1976;27:321–48.

151 13. Endo M, Nakamura S, Araki T, Mochizuki N, Nagatani A. Phytochrome B in the
152 mesophyll delays flowering by suppressing *FLOWERING LOCUS T* expression in
153 *Arabidopsis* vascular bundles. Plant Cell 2005;17:1941-52.

154

155 **Figure legends**

156 **Figure 1. Flowering phenotype and protein stabilization of 35S::PHL-YFP.**

157 (A, B), Plants were grown under 16h light/8h dark long day and 8h light/16h dark short
158 day conditions at 22°C. Mean \pm SD (n \geq 12). (C), 35S::PHL-YFP were grown under
159 continuous white light (cW), continuous dark (cD) for 7 days. Seedlings grown under
160 cD were then exposed to white light for 1 to 24 hour (cD + cW). YFP fluorescence was
161 observed under a laser scanning confocal microscope. Bar=50 μ m

162

163 **Figure 2. PHL-GUS stabilization under light and dark conditions.**

164 Ten-day-old PHLpro::PHL-GUS *phl-1* plants grown under continuous white light (cW)
165 and continuous dark (cD) (A), and 1 to 24h exposure of white light to the cD grown
166 seedlings (B). Bars=1 mm




Figure 1






Figure 2